



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

04. 08. 03

REC'D PCT/PTO 31 JAN 2005
PAT/EP 03 / 50359 #2

REC'D 22 SEP 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02078231.4

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 02078231.4
Demande no:

Anmeldetag:
Date of filing: 02.08.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Tibotec Pharmaceuticals Ltd.
Unit 4, Block 4B, Blanchardstown Corporate
Park
Blanchardstown, Dublin 15
IRLANDE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Broadspectrum 2-amino-benzothiazole sulfonamide HIV protease inhibitors

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K31/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

BROADSPECTRUM 2-AMINO-BENZOTHAZOLE SULFONAMIDE HIV
PROTEASE INHIBITORS

5 The present invention relates to 2-amino-benzothiazole sulfonamides, their use as broadspectrum HIV protease inhibitors, processes for their preparation as well as pharmaceutical compositions and diagnostic kits comprising them. The present invention also concerns combinations of the present 2-aminobenzoxazole sulfonamides with another anti-retroviral agent. It further relates to their use in assays as reference compounds or as reagents.

10 The virus causing the acquired immunodeficiency syndrome (AIDS) is known by different names, including T-lymphocyte virus III (HTLV-III) or lymphadenopathy-associated virus (LAV) or AIDS-related virus (ARV) or human immunodeficiency virus (HIV). Up until now, two distinct families have been identified, i.e. HIV-1 and
15 HIV-2. Hereinafter, HIV will be used to generically denote these viruses.

One of the critical pathways in a retroviral life cycle is the processing of polyprotein precursors by aspartic protease. For instance, with the HIV virus the *gag-pol* protein is processed by HIV protease. The correct processing of the precursor polyproteins by
20 the aspartic protease is required for the assembly of infectious virions, thus making the aspartic protease an attractive target for antiviral therapy. In particular for HIV treatment, the HIV protease is an attractive target.

HIV protease inhibitors (PIs) are commonly administered to AIDS patients in
25 combination with other anti-HIV compounds such as, for instance nucleoside reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), nucleotide reverse transcriptase inhibitors (NtRTIs) or other protease inhibitors. Despite the fact that these antiretrovirals are very useful, they have a common limitation, namely, the targeted enzymes in the HIV virus are able to mutate
30 in such a way that the known drugs become less effective, or even ineffective against these mutant HIV viruses. Or, in other words, the HIV virus creates an ever-increasing resistance against the available drugs.

Resistance of retroviruses, and in particular the HIV virus, against inhibitors is a major
35 cause of therapy failure. For instance, half of the patients receiving anti-HIV combination therapy do not respond fully to the treatment, mainly because of resistance of the virus to one or more drugs used. Moreover, it has been shown that resistant virus is carried over to newly infected individuals, resulting in severely limited therapy options for these drug-naïve patients. Therefore, there is a need in the art for new

compounds for retrovirus therapy, more particularly for AIDS therapy. The need in the art is particularly acute for compounds that are active not only on wild type HIV virus, but also on the increasingly more common resistant HIV viruses.

5 Known antiretrovirals, often administered in a combination therapy regimen, will eventually cause resistance as stated above. This often may force the physician to boost the plasma levels of the active drugs in order for said antiretrovirals to regain
10 effectivity against the mutated HIV viruses. The consequence of which is a highly undesirable increase in pill burden. Boosting plasma levels may also lead to an increased risk of non-compliance with the prescribed therapy. Thus, it is not only
15 important to have compounds showing activity for a wide range of HIV mutants, it is also important that there is little or no variance in the ratio between activity against mutant HIV virus and activity against wild type HIV virus (also defined as fold resistance or FR) over a broad range of mutant HIV strains. As such, a patient may remain on the same combination therapy regimen for a longer period of time since the
20 chance that a mutant HIV virus will be sensitive to the active ingredients will be increased.


25 Finding compounds with a high potency on the wild type and on a wide variety of mutants is also of importance since the pill burden can be reduced if therapeutic levels are kept to a minimum. One additional way of reducing this pill burden is finding anti-HIV compounds with good bioavailability, i.e. a favorable pharmacokinetic and metabolic profile, such that the daily dose can be minimized and consequently also the number of pills to be taken.

Another important characteristic of a good anti-HIV compound is that plasma protein binding of the inhibitor has minimal or even no effect on its potency.


30 Thus, there is a high medical need for protease inhibitors that are able to combat a broad spectrum of mutants of the HIV virus with little variance in fold resistance, have a good bioavailability and experience little or no effect on their potency due to plasma protein binding.

Up until now, several protease inhibitors are on the market or are being developed.
35 One particular core structure (depicted below) has been disclosed in a number of references, such as, WO 95/06030, WO 96/22287, WO 96/28418, WO 96/28463, WO 96/28464, WO 96/28465 and WO 97/18205. The compounds disclosed therein are described as retroviral protease inhibitors.


-3-



WO 99/67254 discloses 4-substituted-phenyl sulfonamides capable of inhibiting multi-drug resistant retroviral proteases.



- 5 Surprisingly, the 2-amino-benzothiazole sulfonamides of the present invention are found to have a favorable virological profile. Not only are they active against wild-type HIV virus, but they also show a broadspectrum activity against various mutant HIV viruses exhibiting resistance against known protease inhibitors.
- 10 The present invention concerns the use of 2-amino-benzothiazole protease inhibitors, having the formula



and *N*-oxides, salts, stereoisomeric forms, racemic mixtures, prodrugs, esters and metabolites thereof, wherein

- 15 R_1 is hexahydrofuro[2,3-*b*]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C_{1-6} alkyl, hydroxy, amino, halogen, amino C_{1-4} alkyl and mono- or di(C_{1-4} alkyl)amino;
- R_2 is hydrogen or C_{1-6} alkyl;
- 20 L is a direct bond, -O-, C_{1-6} alkanediyl-O- or -O- C_{1-6} alkanediyl;
- R_3 is phenyl C_{1-4} alkyl;
- R_4 is C_{1-6} alkyl;
- R_5 is hydrogen or C_{1-6} alkyl;
- R_6 is hydrogen or C_{1-6} alkyl;
- 25 in the manufacture of a medicament useful for inhibiting mutant HIV protease in a mammal infected with said mutant HIV protease. Said mammal in particular is a

human being. The compounds of the present invention are in particular useful in the manufacture of a medicament useful for inhibiting a broad range of mutant HIV proteases.

- 5 The standard of "sensitivity" or alternatively "resistance" of a HIV protease enzyme to a drug is set by the commercially available HIV protease inhibitors. As explained hereinabove, existing commercial HIV protease inhibitors may loose effectivity over time against a population of HIV virus in a patient. The reason being that under pressure of the presence of a particular HIV protease inhibitor, the existing population of HIV virus, usually mainly wild type HIV protease enzyme, mutates into different mutants which a far less sensitive to that same HIV protease inhibitor. If this phenomenon occurs, one talks about resistant mutants. If those mutants are not only resistant to that one particular HIV protease inhibitor, but also to multiple other commercially available HIV protease inhibitors, one talks about multi-drug resistant HIV protease. One way of expressing the resistance of a mutant to a particular HIV protease inhibitor is making the ratio between the EC_{50} of said HIV protease inhibitor against mutant HIV protease over EC_{50} of said HIV protease inhibitor against wild type HIV protease. Said ratio is also called fold resistance (FR).
- 10
- 15
- 20 Many of the mutants occurring in the clinic have a fold resistance of 100 or more against the commercially available HIV protease inhibitors, like saquinavir, indinavir, ritonavir and nelfinavir. Clinically relevant mutants of the HIV protease enzyme can be characterized by a mutation at codon position 10, 71 and/or 84. Examples of such clinical relevant mutant HIV proteases are listed in Table 1.
- 25
- 30 Surprisingly, the compounds of the present invention show a fold resistance ranging between 0.01 and 100 against at least one and in several cases a broad range of clinically relevant mutant HIV proteases. A particular group of compounds of formula (I) are those compounds of formula (I) showing a fold resistance against at least one mutant HIV protease ranging between 0.1 and 100, suitably ranging between 0.1 and 50, and more suitably ranging between 0.1 and 30. Of particular interest are the compounds of formula (I) showing a fold resistance against at least one mutant HIV protease ranging between 0.1 and 20, and even more interesting are those compounds of formula (I) showing a fold resistance against at least one mutant HIV protease ranging between
- 35 0.1 and 10.

Thus, the present invention relates to the use of a compound of formula (I) in the manufacture of a medicament useful for inhibiting replication of a HIV virus having a

mutant HIV protease, in particular a multi-drug resistant mutant HIV protease. It also relates to the use of a compound of formula (I) in the manufacture of a medicament useful for treating or combating a disease associated with HIV viral infection wherein the protease of the HIV virus is mutant, in particular a multi-drug resistant mutant HIV protease.

In other words, the present invention relates to a method of inhibiting a mutant HIV protease, in particular a multi-drug resistant mutant HIV protease, in a mammal infected with said mutant HIV protease, said method comprising contacting said mutant HIV protease in said mammal with an effective amount of a compound of formula (I). The present invention also relates to a method of inhibiting replication of a HIV virus, which has a mutant HIV protease, in particular a multi-drug resistant mutant HIV protease, in a mammal, said method comprising contacting said HIV virus, which has a mutant HIV protease, in said mammal with an effective amount of a compound of formula (I). The present invention further relates to a method of treating or combating a mammalian disease associated with HIV viral infection wherein the protease of the HIV virus is mutant, in particular a multi-drug resistant mutant HIV protease, said method comprising contacting said HIV virus wherein the protease of the HIV virus is mutant infecting said mammal with an effective amount of a compound of formula (I).

A basic nitrogen occurring in the present compounds can be quaternized with any agent known to those of ordinary skill in the art including, for instance, lower alkyl halides, dialkyl sulfates, long chain halides and aralkyl halides.

Whenever the term "substituted" is used in defining the compounds of formula (I), it is meant to indicate that one or more hydrogens on the atom(s) indicated in the expression using "substituted" is replaced with a selection from the indicated group, provided that the indicated atom's normal valency is not exceeded, and that the substitution results in a chemically stable compound, i.e. a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into a therapeutic agent.

As used herein, the term "halo" or "halogen" as a group or part of a group is generic for fluoro, chloro, bromo or iodo.

The term "C₁₋₄alkyl" as a group or part of a group defines straight and branched chained saturated hydrocarbon radicals having from 1 to 4 carbon atoms, such as, for example, methyl, ethyl, propyl, butyl and 2-methyl-propyl.

The term "C₁₋₆alkyl" as a group or part of a group defines straight and branched chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the groups defined for C₁₋₄alkyl and pentyl, hexyl, 2-methylbutyl, 3-methylpentyl and the like.

5 The term "C₁₋₆alkanediyl" as a group or part of a group defines bivalent straight and branched chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, methylene, ethan-1,2-diyl, propan-1,3-diyl, propan-1,2-diyl, butan-1,4-diyl, pentan-1,5-diyl, hexan-1,6-diyl, 2-methylbutan-1,4-diyl, 3-methyl-
10 pentan-1,5-diyl and the like.

As used herein before, the term "one or more" covers the possibility of all the available atoms, where appropriate, to be substituted, preferably, one, two or three.

15 The term "prodrug" as used throughout this text means the pharmacologically acceptable derivatives such as esters, amides and phosphates, such that the resulting *in vivo* biotransformation product of the derivative is the active drug as defined in the compounds of formula (I). The reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8th ed, McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p 13-15) describing prodrugs generally is hereby incorporated. Prodrugs of
20 a compound of the present invention are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of the present invention wherein a hydroxy group, for instance the hydroxy group on the asymmetric carbon atom, or an amino group is bonded to any group that, when the
25 prodrug is administered to a patient, cleaves to form a free hydroxyl or free amino, respectively.

Typical examples of prodrugs are described for instance in WO 99/33795, WO 99/33815, WO 99/33793 and WO 99/33792 all incorporated herein by reference.

30 Prodrugs are characterized by excellent aqueous solubility, increased bioavailability and are readily metabolized into the active inhibitors *in vivo*.

For therapeutic use, the salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically or physiologically acceptable. However, salts having a pharmaceutically unacceptable counterion may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound of formula (I). All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

The pharmaceutically acceptable or physiologically tolerable addition salt forms which the compounds of the present invention are able to form can conveniently be prepared using the appropriate acids, such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; hemisulphuric, nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, aspartic, dodecylsulphuric, heptanoic, hexanoic, nicotinic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methane-sulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pantoic and the like acids.

Conversely said acid addition salt forms can be converted by treatment with an appropriate base into the free base form.

The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt form by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, N-methyl, -D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

Conversely said base addition salt forms can be converted by treatment with an appropriate acid into the free acid form.

The term "salts" also comprises the hydrates and the solvent addition forms that the compounds of the present invention are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

The present compounds may also exist in their tautomeric forms. Such forms, although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

The term stereochemically isomeric forms of compounds of the present invention, as used hereinbefore, defines all possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures which are not interchangeable, which the compounds of the present invention may possess.

Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of all possible stereochemically isomeric forms that said compound might possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically


5 isomeric forms of the compounds of the present invention both in pure form and in admixture with each other are intended to be embraced within the scope of the present invention.

10 Pure stereoisomeric forms of the compounds and intermediates as mentioned herein are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure of said compounds or intermediates. In particular, the term 'stereoisomerically pure' concerns compounds or intermediates having a stereoisomeric excess of at least 80% (i. e. minimum 90% of one isomer and maximum 10% of the other possible isomers) up to a stereoisomeric excess of 100% (i.e. 100% of
15 one isomer and none of the other), more in particular, compounds or intermediates having a stereoisomeric excess of 90% up to 100%, even more in particular having a stereoisomeric excess of 94% up to 100% and most in particular having a stereoisomeric excess of 97% up to 100%. The terms 'enantiomerically pure' and 'diastereomerically pure' should be understood in a similar way, but then having regard to the enantiomeric
20 excess, respectively the diastereomeric excess of the mixture in question.

Pure stereoisomeric forms of the compounds and intermediates of this invention may be obtained by the application of art-known procedures. For instance, enantiomers may be separated from each other by the selective crystallization of their diastereomeric
25 salts with optically active acids or bases. Examples thereof are tartaric acid, dibenzoyltartaric acid, ditoluoyltartaric acid and camphorsulfonic acid. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting
30 materials, provided that the reaction occurs stereospecifically. Preferably, if a specific stereoisomer is desired, said compound would be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

35 The diastereomeric racemates of formula (I) can be obtained separately by conventional methods. Appropriate physical separation methods that may advantageously be employed are, for example, selective crystallization and chromatography, e.g. column chromatography.

It is clear to a person skilled in the art that the compounds of formula (I) contain at least two asymmetric centers and thus may exist as different stereoisomeric forms. These asymmetric centers are indicated with an asterisk (*) in the figure below.



- 5 The absolute configuration of each asymmetric center that may be present in the compounds of formula (I) may be indicated by the stereochemical descriptors R and S, this R and S notation corresponding to the rules described in Pure Appl. Chem. 1976, 45, 11-30. The carbon atom bearing the hydroxy group and marked with the asterisk (*) preferably has the R configuration. The carbon atom bearing the R³ group and
10 marked with the asterisk (*) preferably has the S configuration.

The present invention is also intended to include all isotopes of atoms occurring on the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of
15 hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

Whenever used hereinafter, the term "compounds of formula (I)", or "the present compounds" or similar term is meant to include the compounds of general formula (I), their N-oxides, salts, stereoisomeric forms, racemic mixtures, prodrugs, esters and
20 metabolites, as well as their quaternized nitrogen analogues.


- Some of the compounds of formula (I) have been disclosed in WO 95/06030, i.e.
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid benzyl ester;
25 {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid pyridin-3-ylmethyl ester;
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid thiazol-5-ylmethyl ester;
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
30 3-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;
4-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;
35 5-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;

N-((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-2-methyl-benzamide;

N-((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-4-hydroxy-2-methyl-benzamide;

- 5 N-((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-3-hydroxy-2-methyl-benzamide; and
 ((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-carbamic acid (S)-(tetrahydrofuran-3-yl) ester.

- 10 Hence, the present invention also concerns the compounds of formula (I)



and *N*-oxides, salts, stereoisomeric forms, racemic mixtures, prodrugs, esters and metabolites thereof, wherein

- 15 R_1 is hexahydrofuro[2,3-*b*]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C_{1-6} alkyl, hydroxy, amino, halogen, amino C_{1-4} alkyl and mono- or di(C_{1-4} alkyl)amino;

R_2 is hydrogen or C_{1-6} alkyl;

L is a direct bond, -O-, C_{1-6} alkanediyl-O- or -O- C_{1-6} alkanediyl;

- 20 R_3 is phenyl C_{1-4} alkyl;

R_4 is C_{1-6} alkyl;

R_5 is hydrogen or C_{1-6} alkyl;

R_6 is hydrogen or C_{1-6} alkyl;

provided that the compounds are other than :

- 25 ((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-carbamic acid benzyl ester;
 ((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-carbamic acid pyridin-3-ylmethyl ester;
 ((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-carbamic acid thiazol-5-ylmethyl ester;
 30 ((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-2-(2,6-dimethyl-phenoxy)-acetamide;
 3-amino-((1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl)-2-methyl-benzamide;

- 4-amino-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
 5-amino-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
 5 N-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-methyl-benzamide;
 N-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-4-hydroxy-2-methyl-benzamide;
 N-{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-3-hydroxy-2-methyl-benzamide; and
 10 {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (S)-(tetrahydrofuran-3-yl) ester.

- Interesting compounds are those compounds of formula (I) wherein R¹ is
 15 hexahydrofuro[2,3-b]furanyl or oxazolyl.

- Compounds of particular interest are those compounds wherein -L-R¹ is -O- (hexahydrofuro[2,3-b]furanyl), -O-tetrahydrofuranyl, -O-methyl-(optionally substituted phenyl), -O-methyl-pyridinyl, -O-methyl-thiazolyl, -O-methyl-thiazolyl, -methyl-O-
 20 (optionally substituted phenyl) or optionally substituted phenyl.

- Suitably, one or more of the following restrictions apply to any of the subgroups of the compounds of formula (I) :
- R² is hydrogen;
 25 R³ is phenylmethyl;
 R⁴ is C₁₋₄alkyl, preferably isobutyl;
 R⁵ is hydrogen or methyl;
 R⁶ is hydrogen or methyl.

- A special subgroup of the compounds of formula (I) is defined as the compounds of
 30 formula (I) wherein R⁵ and R⁶ are both hydrogen.

- Compounds of special interest are those compounds of formula (I) wherein R¹ is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected
 35 from C₁₋₆alkyl, hydroxy, amino, chloro, bromo, amino C₁₋₄alkyl and mono-or di(C₁₋₄alkyl)amino;

Preferred compounds are


- {3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid thiazol-5-ylmethyl ester;
-
- 5 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxy-
propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
10 carbamic acid benzyl ester;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid pyridin-3-ylmethyl ester;
- 15 3-amino-N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-3-hydroxy-2-methyl-benzamide;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
20 carbamic acid tetrahydro-furan-3-yl ester;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-methyl-benzamide;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
- 25 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-3-fluoro-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-(4-aminomethyl-2,6-dimethyl-phenoxy)-acetamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
30 propyl}-carbamic acid thiazol-5-ylmethyl ester;
3-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-
sulfonyl)-amino]-propyl}-2-methyl-benzamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-carbamic acid tetrahydro-furan-3-yl ester;
- 35 N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-3-hydroxy-2-methyl-benzamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-(4-iodo-2,6-dimethyl-phenoxy)-acetamide;
-

- 2-(4-aminomethyl-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
 2-(4-amino-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
 5 N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-4-bromo-2-methyl-benzamide;
 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid oxazol-5-ylmethyl ester;
 4-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide; their *N*-oxides, salts and stereoisomeric forms.

Most preferred compounds are those enantiomers having a (1*S*,2*R*)-1-benzyl-2-hydroxy-propyl configuration.

15

Those compounds of formula (I) in a hexahydro-furo[2,3-*b*]furan-3-yl ester form of the carbamic acid derivative occur preferably in a (3*R*,3*aS*,6*aR*) form such as, for instance, {(1*S*,2*R*)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (3*R*,3*aS*,6*aR*)-hexahydro-furo[2,3-*b*]furan-3-yl ester




20

The compounds of formula (I) can generally be prepared using procedures analogous to those procedures described in WO 95/06030, WO 96/22287, WO 96/28418, WO 96/28463, WO 96/28464, WO 96/28465 and WO 97/18205.

- 25 Particular reaction procedures to make the present compounds are described below. In the preparations described below, the reaction products may be isolated from the medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, trituration and chromatography.

Scheme A

-14-



The 2-amino-6-chlorosulfonylbenzothiazole derivative (intermediate a-2) was prepared following the procedure described in EP-A-0,445,926. Intermediates a-4 were prepared by reacting an intermediate a-3, prepared according to the procedure described in WO97/18205 and also depicted in scheme B, with an intermediate a-2 in a reaction-inert solvent such as dichloromethane, and in the presence of a base such as triethylamine and at low temperature, for example at 0°C . The Boc group in the intermediate a-3 is a protective *tert*-butoxycarbonyl group. Another suitable protective group such as phthalimido or benzyloxycarbonyl may conveniently replace it.

Intermediates a-4 may be deprotected with an acid such as hydrochloric acid in isopropanol or with trifluoroacetic acid depending on the nature of the amino group in the 2 position of benzoxazole, in a suitable solvent such as a mixture of ethanol and dioxane, thus preparing an intermediate a-5. Said intermediate a-5 may be further reacted with an intermediate of formula $\text{R}_1\text{-L-C(=O)-OH}$ in the presence of a base such as triethylamine (for alcohols to generate a carbamate) and optionally in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloric acid (EDC) and 1-hydroxybenzotriazole (HOBT) (for carboxylic acids to generate an amide) or an alcohol such as *tert*-butanol, and in a suitable solvent such as dichloromethane; thus forming intermediates a-6.


A convenient way of preparing compounds of formula (I) wherein both R_5 and R_6 are hydrogen can be prepared analogously to the procedure described in scheme A, and whereby one of R_5 or R_6 is replaced by a suitable protective group such as, for example, an acetyl or an alkyloxycarbonyl group. In such a case, deprotection may

-15-

occur simultaneously with the deprotection of the nitrogen atom on the left-hand side of the molecule.

- A number of intermediates and starting materials used in the foregoing preparations are known compounds, while others may be prepared according to art-known methodologies of preparing said or similar compounds.

Scheme B



- Intermediate b-2, corresponding to intermediate a-3 in scheme A, may be prepared by adding an amine of formula $\text{H}_2\text{N-R}_4$ to an intermediate b-1 in a suitable solvent such as isopropanol.

- In scheme B, enantiomerically pure compounds of formula b-2 are only obtained if b-1 is enantiomerically pure. If b-1 is a mixture of stereoisomers, then b-2 will also consist of a mixture of stereoisomers.

- The compounds of formula (I) may also be converted to the corresponding *N*-oxide forms following art-known procedures for converting trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chloro-benzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *tert*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

- The present compounds can thus be used in animals, preferably in mammals, and in particular in humans as pharmaceuticals per se, in mixtures with one another or in the form of pharmaceutical preparations.

Furthermore, the present invention relates to pharmaceutical preparations that as active constituents contain an effective dose of at least one of the compounds of formula (I) in addition to customary pharmaceutically-innocuous-excipients and auxiliaries. The pharmaceutical preparations normally contain 0.1 to 90% by weight of a compound of formula (I). The pharmaceutical preparations can be prepared in a manner known per se to one of skill in the art. For this purpose, at least one of a compound of formula (I), together with one or more solid or liquid pharmaceutical excipients and/or auxiliaries and, if desired, in combination with other pharmaceutical active compounds, are brought into a suitable administration form or dosage form which can then be used as a pharmaceutical in human medicine or veterinary medicine.

Pharmaceuticals which contain a compound according to the invention can be administered orally, parenterally, e.g., intravenously, rectally, by inhalation, or topically, the preferred administration being dependent on the individual case, e.g., the particular course of the disorder to be treated. Oral administration is preferred.

The person skilled in the art is familiar on the basis of his expert knowledge with the auxiliaries that are suitable for the desired pharmaceutical formulation. Beside solvents, gel-forming agents, suppository bases, tablet auxiliaries and other active compound carriers, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, agents for achieving a depot effect, buffer substances or colorants are also useful.

Due to their favorable pharmacological properties, particularly their activity against multi-drug resistant HIV protease enzymes, the compounds of the present invention are useful in the treatment of individuals infected by HIV and for the prophylaxis of these individuals. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses whose existence is mediated by, or depends upon, the protease enzyme. Conditions which may be prevented or treated with the compounds of the present invention, especially conditions associated with HIV and other pathogenic retroviruses, include AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic CNS diseases caused by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis.

Said method of treatment comprises the systemic administration to HIV-infected subjects of an amount effective to combat the conditions associated with HIV virus with multi-drug resistant protease enzyme.

The compounds of the present invention may also find use in inhibiting *ex vivo* samples containing multi-drug resistant HIV-protease or expected to be exposed to multi-drug resistant HIV-protease. Hence, the present compounds may be used to inhibit multi-drug resistant HIV-protease present in a body fluid sample that contains or is suspected to contain or be exposed to multi-drug resistant HIV-protease.

Also, the combination of an antiretroviral compound and a compound of the present invention can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of the present invention, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in treatment of retroviral infections, in particular, in the treatment of infections with multi-drug resistant HIV proteases. Thus, to combat or treat infections with multi-drug resistant HIV protease, or the infection and disease associated with such infections, such as Acquired Immunodeficiency Syndrome (AIDS) or AIDS Related Complex (ARC), the compounds of this invention may be co-administered in combination with for instance, binding inhibitors, such as, for example, dextran sulfate, suramine, polyanions, soluble CD4, PRO-542, BMS-806; fusion inhibitors, such as, for example, T20, T1249, 5-helix, D-peptide ADS-J1; co-receptor binding inhibitors, such as, for example, AMD 3100, AMD-3465, AMD7049, AMD3451 (Bicyclams), TAK 779; SHC-C (SCH351125), SHC-D, PRO-140RT inhibitors, such as, for example, foscarnet and prodrugs; nucleoside RTIs, such as, for example, AZT, 3TC, DDC, DDI, D4T, Abacavir, FTC, DAPD, dOTC, DPC 817; nucleotide RTIs, such as, for example, PMEA, PMPA (tenofovir); NNRTIs, such as, for example, nevirapine, delavirdine, efavirenz, 8 and 9-Cl TIBO (tivicapine), loviride, TMC-125, dapivirine, MKC-442, UC 781, UC 782, Capravirine, DPC 961, DPC963, DPC082, DPC083, calanolide A, SJ-1366, TSAO, 4"-deaminated TSAO, MV150, MV026048; RNase H inhibitors, such as, for example, SP1093V, PD126338; TAT inhibitors, such as, for example, RO-5-3335, K12, K37; integrase inhibitors, such as, for example, L 708906, L 731988, S-1360; protease inhibitors, such as, for example, amprenavir and prodrug GW908, ritonavir, nelfinavir, saquinavir, indinavir, lopinavir, palinavir, BMS 186316, atazanavir, DPC 681, DPC 684, tipranavir, AG1776, mozenavir, GS3333, KNI-413, KNI-272, L754394, L756425, LG-71350, PD161374, PD173606, PD177298, PD178390, PD178392, PNU 140135, TMC114 maslinic acid, U-140690; glycosylation inhibitors, such as, for example, castanospermine, deoxynojirimycin.

The combination may in some cases provide a synergistic effect, whereby viral infectivity and its associated symptoms may be prevented, substantially reduced, or eliminated completely.

5 The compounds of the present invention may also be administered in combination with immunomodulators (e.g., bropirimine, anti-human alpha interferon antibody, IL-2, methionine enkephalin, interferon alpha, HE-2000 and naltrexone) with antibiotics (e.g., pentamidine isothiorate) cytokines (e.g. Th2), modulators of cytokines, chemokines or the receptors thereof (e.g. CCR5) or hormones (e.g. growth hormone) to
10 ameliorate, combat, or eliminate HIV infection and its symptoms. Such combination therapy in different formulations may be administered simultaneously, separately or sequentially. Alternatively, such combination may be administered as a single formulation, whereby the active ingredients are released from the formulation simultaneously or separately.

15 The compounds of the present invention may also be administered in combination with modulators of the metabolism following application of the drug to an individual. These modulators include compounds that interfere with the metabolism at cytochromes, such as cytochrome P450. Some modulators inhibit cytochrome P450. It
20 is known that several isoenzymes exist of cytochrome P450, one of which is cytochrome P450 3A4. Ritonavir is an example of a modulator of metabolism via cytochrome P450. Such combination therapy in different formulations may be administered simultaneously, separately or sequentially. Alternatively, such combination may be administered as a single formulation, whereby the active
25 ingredients are released from the formulation simultaneously or separately. Such modulator may be administered at the same or different ratio as the compound of the present invention. Preferably, the weight ratio of such modulator vis-à-vis the compound of the present invention (modulator:compound of the present invention) is 1:1 or lower, more preferable the ratio is 1:3 or lower, suitably the ratio is 1:10 or
30 lower, more suitably the ratio is 1:30 or lower.

For an oral administration form, compounds of the present invention are mixed with suitable additives, such as excipients, stabilizers or inert diluents, and brought by means of the customary methods into the suitable administration forms, such as tablets, coated
35 tablets, hard capsules, aqueous, alcoholic, or oily solutions. Examples of suitable inert carriers are gum arabic, magnesia, magnesium carbonate, potassium phosphate, lactose, glucose, or starch, in particular, cornstarch. In this case the preparation can be carried out both as dry and as moist granules. Suitable oily excipients or solvents are vegetable

or animal oils, such as sunflower oil or cod liver oil. Suitable solvents for aqueous or alcoholic solutions are water, ethanol, sugar solutions, or mixtures thereof.

Polyethylene glycols and polypropylene glycols are also useful as further auxiliaries for other administration forms.

5

For subcutaneous or intravenous administration, the active compounds, if desired with the substances customary therefore such as solubilizers, emulsifiers or further auxiliaries, are brought into solution, suspension, or emulsion. The compounds of formula (I) can also be lyophilized and the lyophilizates obtained used, for example, for the production of injection or infusion preparations. Suitable solvents are, for example, water, physiological saline solution or alcohols, e.g. ethanol, propanol, glycerol, in addition also sugar solutions such as glucose or mannitol solutions, or alternatively mixtures of the various solvents mentioned.

15 Suitable pharmaceutical formulations for administration in the form of aerosols or sprays are, for example, solutions, suspensions or emulsions of the compounds of formula (I) or their physiologically tolerable salts in a pharmaceutically acceptable solvent, such as ethanol or water, or a mixture of such solvents. If required, the formulation can also additionally contain other pharmaceutical auxiliaries such as surfactants, emulsifiers and stabilizers as well as a propellant. Such a preparation customarily contains the active compound in a concentration from approximately 0.1 to 50%, in particular from approximately 0.3 to 3% by weight.

25 In order to enhance the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions, it can be advantageous to employ α -, β - or γ -cyclodextrins or their derivatives. Also co-solvents such as alcohols may improve the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions. In the preparation of aqueous compositions, addition salts of the subject compounds are obviously more suitable due to their increased water solubility.

30

Appropriate cyclodextrins are α -, β - or γ -cyclodextrins (CDs) or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C_{1-6} alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxy C_{1-6} alkyl, particularly hydroxyethyl, hydroxypropyl or hydroxybutyl; carboxy C_{1-6} alkyl, particularly carboxymethyl or carboxyethyl; C_{1-6} alkyl-carbonyl, particularly acetyl; C_{1-6} alkyloxycarbonyl C_{1-6} alkyl or carboxy C_{1-6} alkyloxy C_{1-6} alkyl, particularly carboxymethoxypropyl or carboxyethoxypropyl; C_{1-6} alkylcarbonyloxy C_{1-6} alkyl, particularly 2-acetyloxypropyl. Especially noteworthy as complexants and/or solubilizers are β -CD, randomly methylated β -CD,

35

2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- γ -CD, 2-hydroxypropyl- γ -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD (2-HP- β -CD).

- 5 The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

- 10 An interesting way of formulating the present compounds in combination with a cyclodextrin or a derivative thereof has been described in EP-A-721,331. Although the formulations described therein are with antifungal active ingredients, they are equally interesting for formulating the compounds of the present invention. The formulations described therein are particularly suitable for oral administration and comprise an antifungal as active ingredient, a sufficient amount of a cyclodextrin or a derivative
- 15 thereof as a solubilizer, an aqueous acidic medium as bulk liquid carrier and an alcoholic co-solvent that greatly simplifies the preparation of the composition. Said formulations may also be rendered more palatable by adding pharmaceutically acceptable sweeteners and/or flavors.

- 20 Other convenient ways to enhance the solubility of the compounds of the present invention in pharmaceutical compositions are described in WO 94/05263, WO 98/42318, EP-A-499,299 and WO 97/44014, all incorporated herein by reference.

- 25 More in particular, the present compounds may be formulated in a pharmaceutical composition comprising a therapeutically effective amount of particles consisting of a solid dispersion comprising (a) a compound of formula (I), and (b) one or more pharmaceutically acceptable water-soluble polymers.

- 30 The term "a solid dispersion" defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, wherein one component is dispersed more or less evenly throughout the other component or components. When said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermodynamics, such a solid dispersion is referred to as "a solid solution". Solid solutions are
- 35 preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered.

The term "a solid dispersion" also comprises dispersions that are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically

-21-

uniform throughout or comprise more than one phase.

5 The water-soluble polymer in the particles is conveniently a polymer that has an apparent viscosity of 1 to 100 mPa.s when dissolved in a 2 % aqueous solution at 20°C solution.

Preferred water-soluble polymers are hydroxypropyl methylcelluloses or HPMC. HPMC having a methoxy degree of substitution from about 0.8 to about 2.5 and a hydroxypropyl molar substitution from about 0.05 to about 3.0 is generally water
10 soluble. Methoxy degree of substitution refers to the average number of methyl ether groups present per anhydroglucose unit of the cellulose molecule. Hydroxy-propyl molar substitution refers to the average number of moles of propylene oxide that have reacted with each anhydroglucose unit of the cellulose molecule.

15 First preparing a solid dispersion of the components, and then optionally grinding or milling that dispersion allows one to prepare the particles as defined hereinabove. Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation, melt-extrusion being preferred.

20 It may further be convenient to formulate the present compounds in the form of nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm. Useful surface modifiers are believed to include those that physically adhere to the surface of the antiretroviral agent but do not chemically bond to the antiretroviral agent.

25 Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

30 Yet another interesting way of formulating the present compounds involves a pharmaceutical composition whereby the present compounds are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition with good bioavailability which can conveniently be manufactured and
35 which is suitable for preparing pharmaceutical dosage forms for oral administration.

Said beads comprise (a) a central, rounded or spherical core, (b) a coating film of a hydrophilic polymer and an antiretroviral agent and (c) a seal-coating polymer layer.

-22-

Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.

5

The route of administration may depend on the condition of the subject, co-medication and the like.

Another aspect of the present invention concerns a kit or container comprising a compound of formula (I) in an amount effective for use as a standard or reagent in a
10 test or assay for determining the ability of a potential pharmaceutical to inhibit multi-drug resistant HIV protease, HIV growth, or both. This aspect of the invention may find its use in pharmaceutical research programs.

15

The compounds of the present invention can be used in phenotypic resistance monitoring assays, such as known recombinant assays, in the clinical management of resistance developing diseases such as HIV. A particularly useful resistance monitoring system is a recombinant assay known as the AntivirogramTM. The AntivirogramTM is a highly automated, high throughput, second generation, recombinant assay that can measure susceptibility, especially viral susceptibility, to the
20 compounds of the present invention. (Hertogs K, de Bethune MP, Miller V *et al.* *Antimicrob Agents Chemother*, 1998; 42(2):269-276, incorporated by reference).

25

Interestingly, the compounds of the present invention may comprise chemically reactive moieties capable of forming covalent bonds to localized sites such that said compound have increased tissue retention and half-lives. The term "chemically reactive group" as used herein refers to chemical groups capable of forming a covalent bond. Reactive groups will generally be stable in an aqueous environment and will usually be carboxy, phosphoryl, or convenient acyl group, either as an ester or a mixed anhydride, or an imidate, or a maleimide thereby capable of forming a covalent bond with
30 functionalities such as an amino group, a hydroxy or a thiol at the target site on for example blood components such as albumine. The compounds of the present invention may be linked to maleimide or derivatives thereof to form conjugates.

35

The dose of the present compounds or of the physiologically tolerable salt(s) thereof to be administered depends on the individual case and, as customary, is to be adapted to the conditions of the individual case for an optimum effect. Thus it depends, of course, on the frequency of administration and on the potency and duration of action of the compounds employed in each case for therapy or prophylaxis, but also on the nature

-23-

- and severity of the infection and symptoms, and on the sex, age, weight, co-medication, and individual responsiveness of the human or animal to be treated and on whether the therapy is acute or prophylactic. Customarily, the daily dose of a compound of formula (I) in the case of administration to a patient approximately 75 kg in weight is 1 mg to 3g, suitably 1 mg to 1g, preferably 3 mg to 0.5 g, more preferably 5 mg to 300 mg. The dose can be administered in the form of an individual dose, or divided into several, e.g. two, three, or four, individual doses.

Experimental Part

10 Preparation of the compounds of formula (I)

The nomenclature used throughout the description is based on Chemical Abstracts Services Nomenclature.

Example 1 : Compound 2

- 15 To a mixture of 825 mg 2-amino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide and 373 mg triethylamine in dichloromethane was added 452 mg 1-[[[(3R,3aS,6aR)-hexahydrofuro[2,3-b]furan-3-yl]oxy]carbonyl]oxy]-2,5-pyrrolidine-dione (described in W09967417). This mixture was stirred at room temperature for 12 hours. After evaporation of dichloromethane under reduced pressure, 20 the crude product was purified on silica, yielding 270 mg 24.8 % compound 2.

Example 2 : Compound 4

- To a mixture of 350 mg 2-methylamino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide and 200 mg triethylamine in dichloromethane 25 was added 210 mg 1-[[[hexahydrofuro[2,3-b]furan-3-yl]oxy]carbonyl]oxy]-2,5-pyrrolidinedione (described in W09967417). This mixture was stirred at room temperature for 12 hours. After evaporation of dichloromethane under reduced pressure, the crude product was purified on silica, yielding 260 mg (55 %) of compound 4.

30 Example 3 : Compound 6

- To a mixture of 420 mg 2-dimethylamino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide and 98 mg triethylamine in dichloromethane was added 230 mg 1-[[[hexahydrofuro[2,3-b]furan-3-yl]oxy]carbonyl]oxy]-2,5-pyrrolidinedione (described in W09967417). This mixture was stirred at room 35 temperature for 12 hours. After evaporation of dichloromethane under reduced pressure, the crude product was purified on silica, yielding 500 mg 90 % of compound 6.

Example 4 : Compound 17

-24-

A mixture of 800 mg of 2-amino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide, 50 mg of HOBT (hydroxybenzotriazol), 420 mg of EDC and 668 mg of (3,4,5-trimethyl-benzyl)-carbamic acid tert-butyl ester compound with hydroxy acetic acid in 80 ml of dichloromethane, was stirred overnight at room

- 5 temperature. The reaction mixture was then washed with water and brine. The organic layer was separated, dried and the solvent evaporated. The residue was purified by column chromatography, yielding 1 g (75 %) of [4-({3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropylcarbonyl}-methoxy)-3,5-dimethyl-benzyl]-carbamic acid tert-butyl ester. This intermediate (500 mg) was further dissolved
10 in methanol (20 ml) and 10 ml of a solution of HCl in isopropanol (5 to 6 N) was added dropwise. The mixture was stirred overnight at room temperature. The solvent was evaporated and the residue was purified on silica yielding 190 mg of compound 17 (43%).

15 Example 5 : Compound 27

- A mixture of 134 mg of 2-methylamino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide, 4 mg of HOBT (hydroxybenzotriazol), 66 mg of EDC and 63 mg of 4-bromo-2-methyl benzoic acid in dichloromethane, was stirred overnight at room temperature. The reaction mixture was then washed with water and
20 brine. The organic layer was separated, dried and the solvent evaporated. The residue was purified by preparative HPLC yielding 25 mg (13 %) of compound 27.

Example 6 : Compound 28

- To a mixture of 4.48 g 2-methylamino-benzothiazole-6-sulfonic acid (3-amino-2-hydroxy-4-phenyl-butyl)-isobutyl amide and 2.73 g triethylamine in dichloromethane
25 was added 3.45 g carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester oxazol-5-ylmethyl ester. This mixture was stirred at room temperature for 12 hours. After evaporation of dichloromethane under reduced pressure, the crude product was purified on silica, yielding 1.02 g 19 % compound 28.

30

The compounds in Table 1, not intended to be limiting the scope of the present invention, have been prepared and tested in support of the presently claimed invention :

Table 1

Name	Number
{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester	1
{{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (3R,3aS,6aR)-hexahydro-furo[2,3-b]furan-3-yl ester	2

{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid thiazol-5-ylmethyl ester	3
{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester	4
{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid (3R,3aS,6aR)-(hexahydro-furo[2,3-b]furan-3-yl) ester	5
{(1S,2R)-1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxy-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester	6
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid benzyl ester	7
N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-(2,6-dimethyl-phenoxy)-acetamide	8
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid pyridin-3-ylmethyl ester	9
3-amino-N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide	10
N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-3-hydroxy-2-methyl-benzamide	11
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid tetrahydro-furan-3-yl ester	12
{(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (S) (tetrahydro-furan-3-yl) ester	13
N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide	14
N-[(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl]-2-(2,6-dimethyl-phenoxy)-acetamide;	15
N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-3-fluoro-2-methyl-benzamide	16
N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-(4-aminomethyl-2,6-dimethyl-phenoxy)-acetamide	17
{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid thiazol-5-ylmethyl ester	18
{(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid thiazol-5-ylmethyl ester trifluoroacetate	19
3-amino-N-[(1S,2R)-1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl]-2-methyl-benzamide trifluoroacetate	20

-27-

- 90%, respectively, of the cells from the cytopathogenic effect of the virus. The toxicity of the compound was measured on the mock-infected cells and was expressed as CC_{50} , which represents the concentration of compound required to inhibit the growth of the cells by 50%. The selectivity index (SI) (ratio CC_{50}/EC_{50}) is an indication of the selectivity of the anti-HIV activity of the inhibitor. Wherever results are reported as e.g. pEC_{50} or pCC_{50} values, the result is expressed as the negative logarithm of the result expressed as EC_{50} or CC_{50} respectively.

Antiviral spectrum:

- Because of the increasing emergence of drug resistant HIV strains, the present compounds were tested for their potency against clinically isolated HIV strains harboring several mutations (Table 2 and 3). These mutations are associated with resistance to protease inhibitors and result in viruses that show various degrees of phenotypic cross-resistance to the currently commercially available drugs such as for instance saquinavir, ritonavir, nelfinavir, indinavir and amprenavir.

Table 2 List with a representative selection of mutant HIV strains (A to F).

Strain	Mutations in HIV protease gene
A	V003I, L010I, V032T, L033M, E035D, S037Y, S037D, M046I, R057R/K, Q058E, L063P, K070T, A071V, I072V, I084V, L089V
B	V003I, L010I, K020R, E035D, M036I, S037N, Q058E, I062V, L063P, A071V, I072M, G073S, V077I, I084V, I085V, L090M
C	V003I, L010I, I015V, L019I, K020M, S037N, R041K, I054V, Q058E, L063P, A071V, I084V, L090M, I093L
D	V003I, L010L/I, I013V, L033I, E035D, M036I, M046I, K055R, R057K, L063P, I066F, A071V, I084V, N088D, L090M
E	V003I, L010I, V011I, A022V, L024I, E035D, M036I, S037T, R041K, I054V, I062V, L063P, A071V, I084V
F	L010F, M046I, M071V, I084V

20 Results:

- As a measure of the broad spectrum activity of the present compounds, Table 3 shows the results of the antiviral testing in terms of pEC_{50} ($= -\log$ of EC_{50}). The fold resistance (FR), defined as $FR = EC_{50}(\text{mutant strain})/EC_{50}(\text{HIV-1 strain LAI})$ is listed in Table 4. For most of the compounds the fold resistance ranges between 0.1 and 100.
- 25 Thus, the present compounds are potent inhibitors of a broad range of mutant strains.

-28-

The toxicity (Tox) is expressed as the pCC_{50} value as determined with mock transfected cells while the pEC_{50} for the wild type is displayed in column WT.

Table 3. Results of the toxicity testing and the resistance testing against strain A to F

5 (expressed as pEC_{50}).

Compound	Strain A	Strain B	Strain C	Strain D	Strain E	Strain F	Tox	WT
1	8.53	8.44	8.38	8.65	8.51	ND	4.16	8.26
2	8.68	8.59	8.54	8.69	8.50	8.45	4.07	8.18
3	7.52	8.05	7.81	7.44	7.66	7.27	4.13	8.34
4	8.44	8.93	8.93	8.93	8.89	8.06	<5	9.34
5	9.37	9.57	9.71	ND	ND	8.71	4.15	9.26
6	6.70	6.89	7.47	6.96	ND	6.15	<4	8.4
7	6.37	7.53	7.49	6.93	7.36	6.11	4.33	8.23
8	7.5	7.87	7.59	7.47	7.56	6.85	<5	8.18
9	6.58	8.25	5.31	7.38	7.62	ND	4.29	8.31
10	7.07	8.03	7.80	7.64	7.88	7.06	4.14	8.04
11	6.95	8.14	8.12	8.08	8.14	6.99	4.24	7.84
12	6.64	8.12	6.72	7.58	8.11	ND	<4	8.37
13	7.39	8.24	8.42	8.13	8.57	6.98	<4	8.52
14	6.05	7.57	6.75	7.40	7.52	ND	4.33	8.42
15	7.29	7.54	7.40	7.30	7.44	6.64	4.04	7.95
16	<4	<4	<4	<4	<4	<4	4.95	5.85
17	7.50	8.18	7.91	7.63	8.12	6.80	4.2	8.15
18	7.51	8.21	8.13	7.67	8.04	6.83	4.07	8.72
19	7.36	7.80	7.88	7.43	7.90	6.80	ND	8.51
20	6.50	7.61	7.40	7.38	7.59	6.12	<4	8.32
22	6.93	7.83	8.19	7.78	8.36	6.1	<4	8.84
23	6.54	8.02	8.06	7.67	8.14	5.20	4.16	8.34
25	7.36	7.70	7.75	7.39	7.76	6.32	4.85	8.39
26	7.52	8.40	8.14	8.08	8.21	7.25	ND	8.57
27	6.80	7.69	5.30	7.07	7.51	6.19	<4	7.72
28	7.71	8.25	8.21	7.43	8.15	7.11	ND	8.60

ND means not determined


Table 4 : fold resistance

-29-

Compound	Strain A	Strain B	Strain C	Strain D	Strain E	Strain F
1	0.5	0.7	0.8	0.4	0.6	-
2	0.3	0.4	0.4	0.3	0.5	0.5
3	6.6	1.9	3.4	7.9	4.8	11.7
4	7.9	2.6	2.6	2.6	2.8	19.1
5	0.8	0.5	0.4	-	-	3.5
6	50.1	32.4	8.5	27.5	-	177.8
7	72.4	5.0	5.5	20.0	7.4	131.8
8	4.8	2.0	3.9	5.1	4.2	21.4
9	53.7	1.1	1000.0	8.5	4.9	-
10	9.3	1.0	1.7	2.5	1.4	9.5
11	7.8	0.5	0.5	0.6	0.5	7.1
12	53.7	1.8	44.7	6.2	1.8	-
13	13.5	1.9	1.3	2.5	0.9	34.7
14	234.4	7.1	48.8	10.5	7.9	-
15	4.6	2.6	3.5	4.5	3.2	20.4
17	4.5	0.9	1.7	3.3	1.1	22.4
18	16.2	3.2	3.9	11.2	4.8	77.6
19	14.1	5.1	4.3	12.0	4.1	51.3
20	66.1	5.1	8.3	8.7	5.4	158.5
22	81.3	10.2	4.5	11.5	3.0	549.5
23	63.1	2.1	1.9	4.7	1.6	1380.4
25	10.7	4.9	4.4	10.0	4.3	117.5
26	11.2	1.5	2.7	3.1	2.3	20.9
27	8.3	1.1	263.0	4.5	1.6	33.9
28	7.8	2.2	2.5	14.8	2.8	30.9

CLAIMS

1. The use of a 2-amino-benzothiazole, having the formula



a *N*-oxide, salt, stereoisomeric form, racemic mixture, prodrug, ester or metabolite

5 thereof, wherein

R_1 is hexahydrofuro[2,3-*b*]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C_{1-6} alkyl, hydroxy, amino, halogen, amino C_{1-4} alkyl and mono- or di(C_{1-4} alkyl)amino;

10 R_2 is hydrogen or C_{1-6} alkyl;

L is a direct bond, -O-, C_{1-6} alkanediyl-O- or -O- C_{1-6} alkanediyl;

R_3 is phenyl C_{1-4} alkyl;

R_4 is C_{1-6} alkyl;

R_5 is hydrogen or C_{1-6} alkyl;

15 R_6 is hydrogen or C_{1-6} alkyl;

in the manufacture of a medicament useful for inhibiting mutant HIV protease in a mammal infected with said mutant HIV protease.

2. The use according to claim 1 wherein

20 R^2 is hydrogen;

R^3 is phenylmethyl;

R^4 is C_{1-4} alkyl, preferably isobutyl;

R^5 is hydrogen or methyl;

R^6 is hydrogen or methyl.

25

3. The use according to claim 1 or 2 wherein both R^5 and R^6 are hydrogen.

4. The use according to any one of claims 1 to 3 wherein - $L-R^1$ is -O-

(hexahydrofuro[2,3-*b*]furanyl), -O-tetrahydrofuranyl, -O-methyl-(optionally substituted

30 phenyl), -O-methyl-pyridinyl, -O-methyl-thiazolyl, -O-methyl-thiazolyl, -methyl-O-

(optionally substituted phenyl) or optionally substituted phenyl.

5. The use according to any one of claims 1 to 4 wherein the compound is

{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-

35 carbamic acid hexahydro-furo[2,3-*b*]furan-3-yl ester;

- {3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid thiazol-5-ylmethyl ester;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
5 {1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxy-
propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid benzyl ester;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
10 hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid pyridin-3-ylmethyl ester;
3-amino-N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-methyl-benzamide;
15 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-3-hydroxy-2-methyl-benzamide;
{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-
carbamic acid tetrahydro-furan-3-yl ester;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
20 hydroxypropyl}-2-methyl-benzamide;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-3-fluoro-2-methyl-benzamide;
25 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-(4-aminomethyl-2,6-dimethyl-phenoxy)-acetamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-carbamic acid thiazol-5-ylmethyl ester;
3-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-
30 sulfonyl)-amino]-propyl}-2-methyl-benzamide;
{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-carbamic acid tetrahydro-furan-3-yl ester;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-
propyl}-3-hydroxy-2-methyl-benzamide;
35 N-{3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-
hydroxypropyl}-2-(4-iodo-2,6-dimethyl-phenoxy)-acetamide;
2-(4-aminomethyl-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-
methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;

2-(4-amino-2,6-dimethyl-phenoxy)-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-acetamide;
N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-4-bromo-2-methyl-benzamide;


- 5 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid oxazol-5-ylmethyl ester;
4-amino-N-{1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-3-hydroxy-2-methyl-benzamide;
or a salt or stereoisomeric form thereof.

10

6. The use according to any one of claims 1 to 5 wherein the mutant HIV protease has a mutation at a position selected from 10, 71 and 84.

7. The use according to any one of claims 1 to 6 wherein the fold resistance of the
15 mutant HIV protease for the compound described in any one of claims 1 to 5 ranges between 0.01 and 100.

8. A compound having the formula



- 20 a *N*-oxide, salt, stereoisomeric form, racemic mixture, prodrug, ester or metabolite thereof, wherein
R₁ is hexahydrofuro[2,3-*b*]furyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently
selected from C₁₋₆alkyl, hydroxy, amino, halogen, aminoC₁₋₄alkyl and mono-or
25 di(C₁₋₄alkyl)amino;
R₂ is hydrogen or C₁₋₆alkyl;
L is a direct bond, -O-, C₁₋₆alkanediyl-O- or -O-C₁₋₆alkanediyl;
R₃ is phenylC₁₋₄alkyl;
R₄ is C₁₋₆alkyl;
30 R₅ is hydrogen or C₁₋₆alkyl;
R₆ is hydrogen or C₁₋₆alkyl;
provided that the compound is other than :
{(1*S*,2*R*)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid benzyl ester;


- {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid pyridin-3-ylmethyl ester;
 {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid thiazol-5-ylmethyl ester;
 5 {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-2-(2,6-dimethyl-phenoxy)-acetamide;
 3-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;
 4-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;
 10 5-amino-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;
 N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-2-methyl-benzamide;
 15 N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-4-hydroxy-2-methyl-benzamide;
 N-[(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl]-3-hydroxy-2-methyl-benzamide; and
 {(1S,2R)-3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid (S)-(tetrahydrofuran-3-yl) ester.
 20
9. A compound according to claim 8 wherein R¹ is hexahydrofuro[2,3-b]furan-3-yl or oxazolyl.
- 25 10. A compound according to claim 8 or 9 having the formula
 {3-[(2-amino-benzothiazole-6-sulfonyl)-isobutyl-amino]-1-benzyl-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
 {1-benzyl-2-hydroxy-3-[isobutyl-(2-methylamino-benzothiazole-6-sulfonyl)-amino]-propyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
 30 {1-benzyl-3-[(2-dimethylamino-benzothiazole-6-sulfonyl)-isobutyl-amino]-2-hydroxypropyl}-carbamic acid hexahydro-furo[2,3-b]furan-3-yl ester;
 or a salt or stereoisomeric form thereof.

ABSTRACT

BROADSPECTRUM 2-AMINO-BENZOTHAZOLE SULFONAMIDE HIV
PROTEASE INHIBITORS

5

The present invention relates to the use of 2-amino-benzothiazoles, having the formula



- wherein R_1 is hexahydrofuro[2,3-b]furanyl, tetrahydrofuranyl, oxazolyl, thiazolyl, pyridinyl, or phenyl optionally substituted with one or more substituents independently selected from C_{1-6} alkyl, hydroxy, amino, halogen, amino C_{1-4} alkyl and mono- or di(C_{1-4} alkyl)amino; R_2 is hydrogen or C_{1-6} alkyl; L is a direct bond, -O-, C_{1-6} alkanediyl-, O- or -O- C_{1-6} alkanediyl; R_3 is phenyl C_{1-4} alkyl; R_4 is C_{1-6} alkyl; R_5 is hydrogen or C_{1-6} alkyl; R_6 is hydrogen or C_{1-6} alkyl; in the manufacture of a medicament useful for inhibiting mutant HIV protease in a mammal infected with said mutant HIV protease.
- 15 It also relates to novel compounds of formula (I).

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.